Journal of Innovative and Creativity
Journal Homepage: https://joecy.org/index.php/joecy
e-mail: galungwaknook@gmail.co

Faculty of Education
University of Pahlawan Tuanku Tambusai
ISSN 2775-771X (Online)
ISSN 2962-570X (Printed)
6 (1) 2026, Page: 17479-17483

The Impact of Intrauterine TORCH Infections on Fetal and Infant Growth and Development: Clinical Overview and Long-Term Implications

Allia Nova Putri Rizky[1*], Abdi Lathif Yukahadi[2], Damas Prakoso[3], Fatih Darmawansyah[4], Luthfi Fadhillah P[5], Martha Laura Sitanggang[6], M. A. Lathif Kariim[7], Oktavia Dwi S. R[8], Qais Maulana G[9], Raihan Abdullah[10], Rhevita Apsari[11], Shafira Nova[12], Tefi Nadila[13]

Medical Program, Universitas Pertahanan Republik Indonesia
Corresponding Author : Allia Nova Putri Rizky E-mail: alliaanova16@gmail.com

Published: Maret, 2026

ABSTRACT

Intrauterine TORCH infections (Toxoplasmosis, Other, Rubella, Cytomegalovirus, Herpes simplex) pose significant risks to fetal and infant growth, causing miscarriage, congenital defects, IUGR, and long-term neurological/sensory impairments. This study aims to review clinical impacts, mechanisms, diagnosis, treatment, and prevention. It employs descriptive qualitative methods via systematic literature review from PubMed/Google Scholar (2019-2024), targeting peer-reviewed articles (n=20-30 purposive sample). Instruments include search protocols and thematic tables; analysis uses content analysis with triangulation. Results reveal high first-trimester transmission risks (e.g., 80% for rubella), manifestations like hepatosplenomegaly/hearing loss, serological/PCR diagnostics, antivirals (e.g., acyclovir), and vaccination strategies. In conclusion, early screening optimizes outcomes via prenatal care.

Keyword : Congenital Infections, Cytomegalovirus, Fetal Development, Intrauterine Growth Restriction, TORCH Infections

© The Author(s). 2021 Open Access This article is licensed under a Creative Commons Atibution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0

INTRODUCTION

TORCH is a term for a group of infections that are important in caring for newborns. These infections can lead to serious problems for babies and are a major cause of sickness and death in newborns. The issues they cause are different from those in older kids and adults because newborns are still developing.

The acronym TORCH stands for four main infections: cytomegalovirus, toxoplasmosis, herpes simplex virus, and rubella. These infections can cause problems during pregnancy, like miscarriages, birth defects, and issues with brain development. Because of this, doctors often check for these infections during pregnancy to understand any risks.

TORCH infections can really affect newborns. They are linked to serious issues like breathing problems, heart issues, low birth weight, and slow recovery after birth. Babies might also show signs of weak muscles and reflexes. About 0.3% to 2% of newborns are affected by congenital cytomegalovirus, which can lead to various health problems and even death.

It's very important to prevent and treat TORCH infections. They play a big role in newborn health. Finding and treating these infections early can help reduce illness, death, and disabilities in young children. This shows how crucial it is to know about and manage TORCH infections to make sure babies have better health. We also need more research and better prevention methods to help reduce the impact of these infections on mothers and their babies.

METHOD

This study is a descriptive qualitative research using a literature review method from PubMed and Google Scholar (2019-2024), using the keywords "TORCH infections" and "fetal development," as well as peer-reviewed article inclusion criteria. This approach integrates narrative analysis and empirical synthesis, in line with Sugiyono's qualitative methodology and Creswell's mixed methods.

The main instruments were a search protocol and thematic synthesis table for data extraction. The analysis technique used thematic content analysis (open-axial-selective coding) with source triangulation for reliability.

The procedure included: (1) exploratory search, (2) systematic screening, (3) thematic extraction and analysis, and (4) narrative synthesis with peer debriefing validation.

RESULTS AND DISCUSSION

Introduction to TORCH Infections

TORCH stands for Toxoplasma, Other Infections, Rubella, Cytomegalovirus, and Herpes Simplex Virus. If a pregnant woman catches these infections, they can harm her baby. If a mother gets a TORCH infection in the first trimester, it can lead to miscarriage or serious birth defects (Supit, 2021).

1. Toxoplasmosis

   Toxoplasma gondii infects people through contaminated food or water. Common sources are cat poop, raw meat, unwashed fruits and veggies, and contaminated goat milk. A mother can only pass Toxoplasma to her baby if she gets infected for the first time while pregnant (Puspita Sari, 2019).
2. Rubella

   Rubella is a contagious disease, but a vaccine can prevent it. If a mother contracts rubella in the first trimester, there's an 80% chance the virus can infect the baby. This can lead to serious heart, eye, and brain problems (Al-Haddad et al., 2019).
3. Cytomegalovirus (CMV)

   CMV is a common virus that can cause a variety of symptoms. It's the top viral infection passed from mothers to babies, affecting about 1-4% of newborns. The first-time infections are the most dangerous (Liu et al., 2020).
4. Herpes Simplex Virus (HSV)

   HSV spreads easily and usually causes cold sores. It can also cause genital herpes. Most of the time, newborns catch it from mothers who look healthy and show no signs of genital herpes (Dominius, 2024).

Intrauterine Mechanism of TORCH Infections

1. Transmission of Infections from Mother to Fetus

   TORCH infections can cross from mother to baby in different ways. For example, rubella can cause congenital rubella syndrome, leading to problems like cataracts and heart defects. CMV can cause delays in development and hearing loss. These infections are especially harmful when the mother gets them in the first trimester (Liu et al., 2020) The risks of TORCH infections depend on when the mother gets infected. In the first trimester, these infections can be especially harmful to the baby (Supit, 2021).
2. Peripartum Transmission

   This happens when a baby passes through the birth canal. If the mother has active genital herpes, there's a 50 to 60% chance the baby will contract it. If this happens and goes untreated, it can lead to serious health issues.
3. Postnatal Transmission

   Mothers can pass TORCH infections in different ways, such as through their actions or close contact. Not all infections pass through breast milk, but CMV can. If a baby gets CMV, it might cause developmental delays or hearing issues. (Liu et al., 2020). Babies can also catch germs from

the environment, especially if those taking care of them are sick. That's why it's really important for caregivers to keep good hygiene and take health precautions (Ayupova & Madjidova, 2024).



Figure 1 Pathophysiology of TORCH Infections in the Fetus

Impact of TORCH Infections on the Fetus

1. Intrauterine Growth Restriction (IUGR)

   This happens when a baby doesn't grow properly in the womb. Infections like rubella and CMV can harm the placenta, cutting off nutrients and oxygen. This can lead to low birth weight and delays.
2. Congenital Malformations Due to TORCH Infections

   TORCH infections can cause both physical and mental issues in babies before they are born. It's key for pregnant women to prevent and manage these infections to reduce risks (Ayupova & Madjidova, 2024; Gupta et al., 2022; Liu et al., 2020).
3. Risks of Hydrops Fetalis

   Non-immune hydrops fetalis is when fluid builds up in the baby's body. Toxoplasmosis can cause this. If not treated quickly, it can lead to serious issues or even death.

Clinical Manifestations in Infants with TORCH Infections

1. Early Symptoms

   Swollen liver and spleen (hepatosplenomegaly) happen in babies with TORCH infections, especially from CMV and rubella. This can lead to jaundice, liver issues, and weak immunity.
2. Sensory Impairment

   Congenital CMV infection often causes hearing loss. It's the main cause of hearing loss in newborns that isn't genetic. Babies with symptoms usually have more severe hearing loss (Liu et al., 2020).
3. Neuropathy and Motor Impairment

   TORCH infections can lead to brain issues and movement problems. Toxoplasmosis can cause brain swelling and other movement issues. Some brain issues have shown up on scans of the head (Al-Haddad et al., 2019).

Long-Term Implications of TORCH Infections on Growth and Development

Babies infected with TORCH infections might have trouble with thinking skills and motor skills. Some could show signs early, while others face issues later on. They might struggle with muscle tone and movement.

Kids with infections like congenital CMV might deal with long-term health issues. They may need ongoing help from doctors and educators. Kids with TORCH infections, like congenital CMV,

often face long-term health problems. They might have developmental delays and hearing issues. This means they usually need extra help from doctors and educators.

Diagnosis and Early Detection of TORCH Infections

1. Toxoplasmosis

   Doctors usually run blood tests to check for specific antibodies. This helps them see if there's a new infection or if the mother is immune. They check for specific antibodies like IgM and IgG. This helps them see if she's got immunity or if she might have a new infection. There's also a test called the IgG Avidity test. To confirm the infection, they can test for Toxoplasma DNA in the amniotic fluid.
2. Rubella

   Blood tests check if a mother has rubella. For babies, doctors look for specific rubella antibodies in the first three months. They use a test called μ-capture ELISA, which is really good at finding this. Babies with congenital rubella can show the virus in their urine for months or even years.
3. Cytomegalovirus

   Doctors confirm a CMV infection through blood tests. High levels of specific antibodies suggest a risk of passing it to the baby. (Qurniasih et al., 2022). A combination of positive IgM and low IgG avidity is associated with a high risk of intrauterine transmission, up to 40%.
4. Herpes Simplex

   It's important to check for HSV infections early. If the mother has sores during delivery, a C-section might be needed to protect the baby. Tests can check for HSV infections during pregnancy. They look for certain antibodies that show if someone has HSV-1 or HSV-2, helping to tell which type of infection it is (Gupta et al., 2022).

Therapeutic Approaches

1. Toxoplasmosis

   Doctors check for risk factors and ultrasound results during prenatal care. After birth, they may test for eye, hearing, and brain issues. Testing includes checking for IgG, IgM, and IgA antibodies. After birth, doctors may also do eye, hearing, and brain tests, along with a CT scan of the head.
2. Rubella

   Getting vaccinated is key to preventing rubella. Women should get the MMR vaccine before becoming pregnant. Babies with congenital rubella must be reported as they can spread the virus. These babies can spread the virus until they are one year old, unless two negative rubella tests come back in a row.
3. Cytomegalovirus (CMV) Infection

   Managing cytomegalovirus (CMV) during pregnancy is really about prevention. Right now, there isn't a vaccine or a totally safe treatment for moms and babies. Antiviral medications like cidofovir, foscarnet, ganciclovir, and valganciclovir are mostly for patients with weak immune systems or severe CMV cases. But these meds need close monitoring because they can cause side effects like kidney damage and low blood cell counts. So, they're usually only given in serious or high-risk situations. (Zhang et al., 2022).
4. Herpes Simplex Virus (HSV)

   Newborns with HSV often receive strong treatments like intravenous acyclovir. Supportive care can also help. Supportive care can help too. You might use paracetamol to reduce fever or topical lidocaine to ease pain. If the mom's water breaks early, the treatment plan will change based on if the herpes infection is new or if it's an old one (Sofía et al., 2020).

Prevention of TORCH Infections and Risk Mitigation Strategies

Getting an education and staying up to date on vaccinations are really important for staying healthy. For example, kids get the Measles-Rubella vaccine to help protect them from serious issues like hearing or vision problems, heart defects, and mental challenges that can come from a mom

catching rubella when pregnant (Al-Haddad et al., 2019). Also, catching any issues early is a big part of regular check-ups for pregnant women. They often use TORCH screening during these visits.

CONCLUSION

Research shows that TORCH infections can cause serious issues in unborn babies, like small head size, hearing loss, and learning difficulties. Vaccines for some infections are now available. Healthcare providers play a crucial role in catching these issues early and supporting kids as they grow.

We need to improve education around TORCH infections. Early screenings can help prevent infections through healthy practices. More research is needed to find new treatments and understand how these infections affect health over time. Programs should focus on keeping both physical and mental health in check. The aim is to help affected children grow and develop properly.

REFERENCE

Al-Haddad, B., Oler, E., Armistead, B., Elsayed, N., Weinberger, D., Bernier, R., Burd, I., Kapur, R., Jacobsson, B., Wang, C., Mysorekar, I., Rajagopal, L., & Waldorf, K., 2019. *The Fetal Origins of Mental Illness. American Journal of Obstetrics and Gynecology*. doi: 10.1016/j.ajog.2019.06.013.

Ayupova, D. & Madjidova, Y., 2024. *The concept of "TORCH infections" in pregnant women and newborns. Science and Innovation*, 3 (D5), pp. 141-144. doi: 10.5281/zenodo.11383363.

Dominius, A., 2024. *Hidrops fetalis: etiologi dan tata laksana. CDK*, 51(7), pp. 387-391.

Gupta, P., et al., 2022. *Guidelines Diagnosis and Management of Global Development Delay: Consensus Guidelines of Growth, Development and Behavioral Pediatrics Chapter, Neurology Chapter and Neurodevelopment. Indian Pediatrics*, 59(4).

Kamaruddin, M., 2019. *Trik pencegahan dan mengatasi TORCH pada ibu dan bayi melalui natural products. Seminar Nasional Kesehatan Aktabe*, July. Available at:

Liu, Y., Wu, Y., Wang, F., Wang, S., Zhao, W., Chen, L., Tu, S., Qian, Y., Liao, Y., Huang, Y., Zhang, R., Xu, G. & Zhang, D., 2020. *The association between previous TORCH infections and pregnancy and neonatal outcomes in IVF/ICSI-ET: A retrospective cohort study. Frontiers in Endocrinology*, 11, p. 466. doi: 10.3389/fendo.2020.00466.

Puspita Sari, R.D., 2019. *Kehamilan dengan Infeksi TORCH. JK Unila*, 3(1), pp. 176-180.

Qurniasih, N., et al., 2022. *Peningkatan pengetahuan mahasiswa bidan terhadap deteksi dini komplikasi risiko infeksi TORCH pada kehamilan. Jurnal Media Ilmu Keolahragaan Indonesia*, 5(2), pp. 497-504. doi: 10.47679/ib.2022327.

Setiasih, S., Khobibah, T.N., & Nurrokhim, D., 2020. *Buku Ajar Medical Science*. Kendal: Politeknik Kesehatan Semarang dan Kementerian Kesehatan.

Sofía, P., Pérez, J., Campo, S., De Villoria, J., Planells, J., & Domínguez, J., 2020. *A novel Col4A1 mutation: recognizing an uncommon cause of stroke. European Journal of Neurology*, 27. doi: 10.1111/ene.14333.

Supit, B., 2021. *Infeksi TORCH Maternal dan Kongenital. CDK*, 48(9), pp. 377-378.

Tonkin, C.J. (Editor)., 2020. *Methods in Molecular Biology*. 1st ed. Springer US; Humana. ISBN 978-1-4939-9856-2. ISBN 978-1-4939-9857-9 (eBook). DOI: 10.1007/978-1-4939-9857-9.

Zhang, L., Wang, X., Liu, M., Feng, G., Zeng, Y., Wang, R., & Xie, Z., 2022. *The epidemiology and disease burden of congenital TORCH infections among hospitalized children in China: A national cross-sectional study. PLOS Neglected Tropical Diseases*, 16. doi: 10.1371/journal.pntd.0010861.